

A COMBINATORIAL STUDY ON QUIVER VARIETIES

SHIGEYUKI FUJII AND SATOSHI MINABE

ABSTRACT. This is an expository paper which has two parts. In the first part, we study quiver varieties of affine A -type from a combinatorial point of view. We present a combinatorial method for obtaining a closed formula for the generating function of Poincaré polynomials of quiver varieties in rank 1 cases. Our main tools are cores and quotients of Young diagrams. In the second part, we give a brief survey of instanton counting in physics, where quiver varieties appear as moduli spaces of instantons. We discuss combinatorial aspects of its recent developments.

CONTENTS

1. Introduction	1
2. Preliminaries on combinatorics	3
3. Quiver varieties and torus actions	10
4. Enumerative geometry on quiver varieties	14
5. Relations to instanton counting	19
Appendix A. Hilbert schemes of points on surfaces	21
References	24

1. INTRODUCTION

Quiver varieties, which were introduced by Nakajima [36], stand at the corner of a beautiful interplay between combinatorics, geometry, representation theory, and mathematical physics. This expository paper has two purposes. The first one is to study quiver varieties of type $A_{l-1}^{(1)}$ from a combinatorial viewpoint (§2-§4). The second is to give a brief survey of recent developments of instanton counting in physics, especially results of Maeda *et al.* [32, 33, 34], and to discuss combinatorial aspects of them (§5).

The goal in the first part is to compute Poincaré polynomials of quiver varieties in rank 1 case via a combinatorial method. Note that Poincaré polynomials of quiver varieties have already been obtained by Nakajima (see Remark 4.11), and our method just confirms his result. However, we hope that our combinatorial viewpoint gives a new insight into the subject. Our strategy is as follows. There is natural torus actions on quiver varieties. Then by localization principle, we can compute topologies of them by studying the fixed point set. We see that they further reduce to enumeration of Young diagrams. More importantly, we compute the Poincaré polynomial of a quiver variety not one by one, but a *generating function* of them. For that purpose, we use some combinatorial machineries, which are *cores* and *quotient* of Young diagrams. They arise from a division algorithm for

2000 *Mathematics Subject Classification.* Primary 14C05, 14D21; Secondary 05A19, 05E10.

Key words and phrases. Young diagram, core, quotient, quiver variety, instanton.

Young diagrams analogous to that for integers and play important roles in our discussions. They also appear in recent studies of supersymmetric gauge theories by Maeda *et al.* [32, 33, 34] (in slightly different terminologies). The purpose of the second part is to explain it from a mathematical viewpoint. We hope that our exposition is also useful for physics oriented readers.

We introduce quiver varieties of type $A_{l-1}^{(1)}$ as follows. Let Γ be a cyclic group of order l . Then Γ acts on the framed moduli space $\mathcal{M}_r(n)$ of torsion free sheaves E on $\mathbb{CP}^2$ with rank r , $c_2 = n$, where the framing is a trivialization at the line at infinity l_∞ . We consider its Γ -fixed point components,

$$\mathcal{M}_r(n)^\Gamma = \bigsqcup_{\substack{\mathbf{v} \\ |\mathbf{v}|=n}} \mathcal{M}_{\mathbf{w}}(\mathbf{v})$$

where $\mathbf{w}$ and $\mathbf{v}$ are isomorphism classes of Γ -modules. These $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$'s are *quiver varieties of type $A_{l-1}^{(1)}$* . Note that this definition, which is taken from [42], is slightly different from the original one. The original definition was based on the so-called ADHM description of instantons on so-called ALE spaces [25].

The space $\mathcal{M}_r(n)$ was also introduced by Nakajima [37], as a resolution of the moduli space of instantons on $\mathbb{R}^4$. Then quiver varieties can be seen as resolutions of the moduli spaces of instantons on $\mathbb{C}^2/\Gamma$. It turns out that generating functions of Euler characteristics of quiver varieties can be identified with the partition functions of certain supersymmetric field theory, which is important in mathematical physics. This is one of our motivations to study quiver varieties. We summarize some basic facts of instanton counting in §5. Here we should mention a beautiful work by physicists Fucito *et al.* [11]. They studied instanton moduli spaces on ALE spaces by a combinatorial method. The present work grew out of an attempt at rigorous understanding of their idea mathematically. In this paper, we give a precise mathematical formulation of the idea in [11], and complete proofs to the results announced there.

Let us explain more detail about our analysis in the first part. There is a natural action of $(r+2)$ -dimensional torus T on a quiver variety $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$, induced from that on $\mathcal{M}_r(n)$. We study their fixed point sets. It can be shown that fixed points on a quiver variety $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ are parametrized by a certain set $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ of Young diagrams, which is determined by the *coloring of Young diagrams* (see §2 for definition). Then localization theorem tells us that we can compute Euler characteristics of quiver varieties via enumeration of $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$. We can also compute Poincaré polynomials by studying tangent spaces at fixed points. As mentioned above, what we want to compute is the generating function of them with respect to $\mathbf{v}$:

$$\mathcal{Z}_{\mathbf{w}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) \stackrel{\text{def.}}{=} \sum_{\substack{m \geq 0 \\ \mathbf{v} \in (\mathbb{Z}_{\geq 0})^l}} b_m(\mathcal{M}_{\mathbf{w}}(\mathbf{v})) \mathbf{t}^m \mathbf{q}^{|\mathbf{v}|} \prod_{i=0}^{l-1} \mathbf{r}_i^{v_i}.$$

The problem is that how to get a closed formula for $\mathcal{Z}_{\mathbf{w}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$. In §4.2, we give a closed formula for $\mathcal{Z}_{\mathbf{w}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$ when $\mathbf{w}$ is a class $\mathbf{e}_j$ of 1-dimensional Γ -module. First, we show a remarkable factorization property of the generating function of the Poincaré polynomials in Theorem 4.6. It decomposes into the product of the contributions of *cores* and those of *quotients* of Young diagrams in $\mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$:

$$\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) = \mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) \mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}}).$$

Next, we compute each part separately. Computations of the core part $\mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}})$ is purely combinatorial. On the other hand, we use geometry of Hilbert scheme of points, to get the quotients part $\mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$. Then we obtain a closed formula for $\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$ in Theorem 4.10, where $\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$ is expressed as a ratio of the Riemann theta function and the Dedekind eta function. This result is due to Nakajima, as we mentioned earlier.

The paper is structured as follows. We devote §2 to combinatorial preliminaries. The most important notions in §2 are *cores* and *quotients* of Young diagrams. In §3, we introduce quiver varieties and torus actions on them. We study their fixed point sets. In §4, we compute two kinds of global invariants on quiver varieties. In §4.1, we consider ‘equivariant volumes’ of quiver varieties, which are defined in §4.1.1. In §4.2, we study generating functions of Poincaré polynomials and Euler characteristics of quiver varieties, by using combinatorics prepared in §2. In §5, we summarize recent developments of instanton counting in physics and discuss their combinatorial aspects. In appendix A, we collect some facts about Hilbert scheme of points which are used in this paper.

Acknowledgements. The authors would like to thank H. Awata, H. Miyachi, W. Nakai, H. Nakajima, T. Nakatsu, M. Namba, Y. Nohara, Y. Hashimoto, Y. Ito, T. Sasaki, Y. Tachikawa, K. Takasaki, and K. Ueda for valuable discussions and comments. The authors express their deep gratitudes to M. Hamanaka, S. Moriyama, and A. Tsuchiya for their advices and warm encouragements, and especially to H. Kanno for suggesting a problem and reading the manuscript carefully. This work was started while the authors enjoyed the hospitality of the Fields Institute at University of Toronto on the fall of 2004. The authors are grateful to K. Hori for invitation. Throughout this work, the authors’ research was supported in part by COE program in mathematics at Nagoya University.

2. PRELIMINARIES ON COMBINATORICS

In this section, we first fix some notations on Young diagrams which are used through the paper. We also introduce the notion of *quotients* and *cores* for Young diagrams. Our basic references in this section are [21, §2.7] and [50, §3] (see also [53, Exercise 7.59] and [30, I.1. Example 8]).


2.1. Notations on Young diagrams.

2.1.1. *Young diagrams.* A partition is a sequence of non-increasing non-negative integers $\lambda = (\lambda_1, \dots, \lambda_m)$. The corresponding Young diagram is a collection of rows of square boxes which are left adjusted and with λ_i boxes in the i th row for $i = 1, \dots, m$. We identify partitions with their corresponding Young diagrams.

Definition 2.1. (i) The *weight* $|Y|$ of a Young diagram Y is the number of boxes in Y . In terms of the partition λ corresponding to Y , $|Y| = \sum_i \lambda_i$.
(ii) The *total weight* $|\underline{Y}|$ of an l -tuple of Young diagrams $\underline{Y} = (Y_1, \dots, Y_l)$ is defined by $|\underline{Y}| \stackrel{\text{def.}}{=} \sum_{k=1, \dots, l} |Y_k|$.

Notation 2.2. (i) Let $\mathcal{P}$ be the set of Young diagrams. Then we have

$$\mathcal{P} = \bigsqcup_{n \geq 0} \mathcal{P}(n),$$

FIGURE 1. Y_α (dotted) and Y_β (solid)

where $\mathcal{P}(n) = \{Y \in \mathcal{P} \mid |Y| = n\}$.

(ii) Let $\mathcal{P}_l$ be the set of l -tuples of Young diagrams. Then we have

$$\mathcal{P}_l = \bigsqcup_{n \geq 0} \mathcal{P}_l(n),$$

where $\mathcal{P}_l(n) = \{\underline{Y} \in \mathcal{P}_l \mid |\underline{Y}| = n\}$.

2.1.2. Functions on Young diagrams. For a Young diagram Y , we define $l_Y(s)$ and $a_Y(s)$ as

$$l_Y(s) \stackrel{\text{def.}}{=} \lambda_h - k, \quad a_Y(s) \stackrel{\text{def.}}{=} \lambda'_k - h,$$

where $s = (h, k) \in (\mathbb{Z}_{>0})^2$, λ_h is the length of the h -th row of Y , and λ'_k is the length of the k -th column of Y .

Definition 2.3. (i) The box $s \in Y$ sitting on the i -th row and the j -th column is called (i, j) -component of Y and denoted by $s = (i, j)$.

(ii) For $s \in Y$, we define the *leg length* by $l(s) \stackrel{\text{def.}}{=} l_Y(s)$, the *arm length* by $a(s) \stackrel{\text{def.}}{=} a_Y(s)$, and the *hook length* by $h(s) \stackrel{\text{def.}}{=} l(s) + a(s) + 1$.

In the later sections, we need the following generalization of the definition of hook length.

Definition 2.4. Take two Young diagrams Y_α and Y_β . For $s \in (\mathbb{Z}_{>0})^2$ we define

$$h_{Y_\alpha, Y_\beta}(s) \stackrel{\text{def.}}{=} l_{Y_\beta}(s) + a_{Y_\alpha}(s) + 1.$$

We call $h_{Y_\alpha, Y_\beta}(s)$ *relative hook length of s with respect to Y_β* when $s \in Y_\alpha$.

Example 2.5. In Figure 1, the Young diagram drawn by dotted lines is Y_α and the Young diagram drawn by solid lines is Y_β . The number of diamonds (resp. stars) is $l_{Y_\beta}(s)$ (resp. $a_{Y_\alpha}(s)$). Then we have $h_{Y_\alpha, Y_\beta}(s) = 7$.

2.1.3. Removing a hook.

Definition 2.6. For $s = (h, k) \in Y$, we define the *hook* of s as

$$H_s(Y) \stackrel{\text{def.}}{=} \{(i, j) \in Y \mid i = h, j \geq k \text{ or } j = k, i > h\}.$$

By definition, $h(s) = \#H_s(Y)$. We call a hook with n boxes a *hook of length n* .

Definition 2.7. For $s = (h, k) \in Y$, we define the *rim* of s as


$$R_s(Y) \stackrel{\text{def.}}{=} \{(i, j) \in Y \mid i = \lambda'_j \text{ or } j = \lambda_i\} \cap \{(i, j) \in Y \mid i \geq h, j \geq k\}.$$

Then we have

$$\#R_s(Y) = \#H_s(Y) = h(s).$$

By definition, removing a hook $H_s(Y)$ from Y means that removing all boxes in $R_s(Y)$ from Y .

Example 2.8. Let Y be a Young diagram given by a partition $(5, 4, 3, 2, 1)$. We remove $H_{(2,2)}(Y)$ from Y as follows:



2.1.4. Coloring of Young diagrams.

Definition 2.9. Let $Y \in \mathcal{P}$. Let us fix an integer $l \geq 2$. Assume that we are given a $\gamma \in \mathbb{Z}/l\mathbb{Z}$. Then the l -residue of a box $s \in Y$ is defined by the following rule: if s is the (i, j) -component of Y , then

$$(1) \quad \text{res}(s) = \gamma - i + j + l\mathbb{Z} \in \mathbb{Z}/l\mathbb{Z}.$$

We call this assignment of an element of $\mathbb{Z}/l\mathbb{Z}$ to each box in Y as γ -coloring of a Young diagram Y .

We extend this coloring to tuples of Young diagrams.

Definition 2.10. Let $\underline{Y} = (Y_1, \dots, Y_r) \in \mathcal{P}_r$. Assume that we are given $\gamma_1, \dots, \gamma_r \in \mathbb{Z}/l\mathbb{Z}$. If a box $s \in \underline{Y}$ is a (i, j) -component of Y_k , then we color s with

$$\text{res}(s) = \gamma_k - i + j + l\mathbb{Z} \in \mathbb{Z}/l\mathbb{Z}.$$

We call this assignment as a $(\gamma_1, \dots, \gamma_r)$ -coloring of an r -tuple of Young diagrams $\underline{Y}$.

Example 2.11. Take $r = 2$, $l = 3$, and $(\gamma_1, \gamma_2) = (0, 2) \in (\mathbb{Z}/3\mathbb{Z})^2$. Let $\underline{Y} = (Y_1, Y_2) \in \mathcal{P}_2$ which is given by a pair of partitions $(\lambda_1, \lambda_2) = ((4, 3, 1), (3, 2, 1, 1))$. Then (γ_1, γ_2) -coloring of $\underline{Y}$ is given as follows:

$$\left(\begin{array}{|c|c|c|c|} \hline 1 & & & \\ \hline 2 & 0 & 1 & \\ \hline 0 & 1 & 2 & 0 \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 2 & & \\ \hline 0 & & \\ \hline 1 & 2 & \\ \hline 2 & 0 & 1 \\ \hline \end{array} \right).$$

2.2. The Maya diagram.

Definition 2.12. A *Maya diagram* is a sequence $\{\mathbf{m}(\nu)\}_{\nu \in \mathbb{Z}}$ which consists of 0 or 1 and satisfies the following property: there exist $N, M \in \mathbb{Z}$ such that for all $\nu > N$ (resp. $\nu < M$), $\mathbf{m}(\nu) = 1$ (resp. $\mathbf{m}(\nu) = 0$).

It is well-known that there is a one-to-one correspondence between the set of Maya diagrams and the set of Young diagrams. We explain how to associate a Young diagram to a Maya diagram. Let us place Young diagrams on the (x, y) -plane by the following way: the bottom-left corner of the Young diagram is at origin $(0, 0)$ and a box in the Young diagram is an unit square. We call an upper-right borderline of $\{x\text{-axis}\} \cup \{y\text{-axis}\} \cup Y$ the *extended borderline* of Y and denote it by ∂Y . The line defined by $y = x$ is called the *medium*.




FIGURE 2. Young diagram and Maya diagram

Example 2.13. (See Figure 2.) Let Y be the Young diagram corresponding to the partition $(3, 2, 1)$. In Figure 2, ∂Y is shown by a bold line.

The Maya diagram $\{\mathbf{m}_Y(\nu)\}_{\nu \in \mathbb{Z}}$ corresponding to a Young diagram Y is defined as follows. We give the direction to the extended borderline ∂Y of Y , which goes from $(0, +\infty)$ to $(+\infty, 0)$ (see Figure 2). Then each edge of the extended borderline is numbered by $\nu \in \mathbb{Z}$, if we set the edge which is located at the next to the medium to be 0. Next we encode a edge $\downarrow$ (resp. $\rightarrow$) to 0 (resp. 1). By this way, we have a 0/1 sequence $\{\mathbf{m}_Y(\nu)\}_{\nu \in \mathbb{Z}}$, where each $\mathbf{m}_Y(\nu)$ corresponds to a edge of ∂Y .

Notation 2.14. For the sake of brevity, we denote an infinite sequence of 0's (resp. 1's) by $\underline{0}$ (resp. $\underline{1}$).

Example 2.15. The Maya diagram for the Young diagram in Figure 2 is

$$\cdots, 0, 0, 1, 0, 1 \mid 0, 1, 0, 1, 1, \cdots.$$

We express this as

$$\underline{0}, 1, 0, 1 \mid 0, 1, 0, \underline{1}.$$

Here $\mid$ represents the position of the medium. This means that the medium's neighbor on the right is the $\mathbf{m}_Y(0)$.

Remark 2.16. Since the number of edges $\rightarrow$ before the medium is same as the number of edges $\downarrow$ after the medium, we have

$$\#\{\mathbf{m}_Y(\nu) = 1 \mid \nu < 0\} = \#\{\mathbf{m}_Y(\nu) = 0 \mid \nu \geq 0\}.$$

Remark 2.17 (Removing a hook in terms of the Maya diagrams). Fix $(x, y) \in Y$ whose hook length is n . Let $Y^{(x,y)}$ be the Young diagram obtained by removing the hook of length n corresponding to (x, y) from Y . Then the Maya diagram $\{\mathbf{m}_{Y^{(x,y)}}(\nu)\}_{\nu \in \mathbb{Z}}$ of $Y^{(x,y)}$ is given by

$$(2) \quad \mathbf{m}_{Y^{(x,y)}}(\nu) = \begin{cases} \mathbf{m}_Y(\nu) & \text{if } \nu \neq x, y \\ \mathbf{m}_Y(y) & \text{if } \nu = x \\ \mathbf{m}_Y(x) & \text{if } \nu = y. \end{cases}$$

Example 2.18. Let us remove the hook of length 5 from the Young diagram Y shown in Figure 2 by means of the Maya diagram of Y . This is equivalent to exchanging $\mathbf{m}_Y(-3) =$

1 and $\mathbf{m}_Y(2) = 0$, shown as follows:

$$\underline{0}, \widehat{1}, 0, 1 \mid 0, 1, \widehat{0}, \underline{1} \longmapsto \underline{0}, \widehat{0}, 0, 1 \mid 0, 1, \widehat{1}, \underline{1},$$

where $\widehat{}$ represents locations of exchanges.

2.3. Quotients and cores for Young diagrams.

2.3.1. *Quotients.* For each $0 \leq i \leq l-1$, we define

$$(3) \quad \mathbf{m}_{Y_i^*}(n) \stackrel{\text{def.}}{=} \mathbf{m}_Y(ln + i), \quad n \in \mathbb{Z}.$$

Then we have an l -tuple $\left(\left\{ \mathbf{m}_{Y_0^*}(n) \right\}_{n \in \mathbb{Z}}, \dots, \left\{ \mathbf{m}_{Y_{l-1}^*}(n) \right\}_{n \in \mathbb{Z}} \right)$ of subsequences $\{\mathbf{m}_Y(\nu)\}_{\nu \in \mathbb{Z}}$. Each $\{\mathbf{m}_{Y_i^*}(n)\}_{n \in \mathbb{Z}}$ is also a Maya diagram.

Definition 2.19. l -quotient for Y is the l -tuple of Young diagrams $\underline{Y}_{(l)}^* \stackrel{\text{def.}}{=} (Y_0^*, \dots, Y_{l-1}^*)$ corresponding to the Maya diagrams $\left(\left\{ \mathbf{m}_{Y_0^*}(n) \right\}_{n \in \mathbb{Z}}, \dots, \left\{ \mathbf{m}_{Y_{l-1}^*}(n) \right\}_{n \in \mathbb{Z}} \right)$.

Remark 2.20. The hook of length nl in Y corresponds to a hook of length n in $\underline{Y}_{(l)}^*$.

For the latter purpose, let us define the following

Definition 2.21. We define $k_i(Y) \in \mathbb{Z}$ ($i = 0, \dots, l-1$) by the following condition:

$$(4) \quad \# \{ \mathbf{m}_{Y_i^*}(n) = 1 \mid n < k_i(Y) \} = \# \{ \mathbf{m}_{Y_i^*}(n) = 1 \mid n \geq k_i(Y) \}.$$

Remark 2.22. (i) Notice that $k_i(Y) \neq 0$ in general, since we take a subsequence. However

$$\sum_{i=0}^{l-1} k_i(Y) = 0.$$

(ii) $(k_0(Y), \dots, k_{l-1}(Y))$ is invariant under removing a hook from Y .

Example 2.23. Let us consider the 5-quotient for the Young diagram Y shown in Figure

2. (i) In this case,

$$\begin{aligned} \{ \mathbf{m}_Y(\nu) \} &= \{ \underline{0}, 0, 0, 1, 0, 1 \mid 0, 1, 0, 1, 1, \underline{1} \}, \\ \{ \mathbf{m}_{Y_0^*}(n) \} &= \{ \underline{0}, 0, \quad \quad \quad 0, \quad \quad \quad \underline{1} \}, \\ \{ \mathbf{m}_{Y_1^*}(n) \} &= \{ \underline{0}, \quad 0, \quad \quad \quad 1, \quad \quad \quad \underline{1} \}, \\ \{ \mathbf{m}_{Y_2^*}(n) \} &= \{ \underline{0}, \quad \quad 1, \quad \quad \quad 0, \quad \quad \quad \underline{1} \}, \\ \{ \mathbf{m}_{Y_3^*}(n) \} &= \{ \underline{0}, \quad \quad \quad 0, \quad \quad \quad 1, \quad \quad \quad \underline{1} \}, \\ \{ \mathbf{m}_{Y_4^*}(n) \} &= \{ \underline{0}, \quad \quad \quad \quad 1, \quad \quad \quad \quad 1, \quad \quad \quad \underline{1} \}. \end{aligned}$$

Therefore the 5-quotient of Y is

$$(Y_0^*, Y_1^*, Y_2^*, Y_3^*, Y_4^*) = (\emptyset, \emptyset, \square, \emptyset, \emptyset).$$

(ii) Let us remove the hook of length 5 from Y by means of 5-quotient $(Y_0^*, \dots, Y_4^*)$ of Y . This is equivalent to exchanging $\mathbf{m}_{Y_2^*}(-1) = 1$ and $\mathbf{m}_{Y_2^*}(0) = 0$, shown as follows:

$$\{ \mathbf{m}_{Y_2^*}(n) \} = \{ \underline{0}, \widehat{1} \mid \widehat{0}, \underline{1} \} \longmapsto \{ \underline{0}, \widehat{0} \mid \widehat{1}, \underline{1} \}.$$

2.3.2. Cores.

Definition 2.24. A Young diagram is called l -core if its l -quotient $\underline{Y}_{(l)}^*$ is empty. We denote $\mathcal{C}^{(l)}$ by the subset of $\mathcal{P}$ which consists of l -cores.

Example 2.25. The set of 2-cores $\mathcal{C}^{(2)}$ consists of the empty partition and ‘stairs-like’ Young diagrams which correspond to partitions $(m-1, m-2, \dots, 1)$ for $m \geq 2$.

Definition 2.26. Let $Y^{(l)}$ be the Young diagram obtained by removing as many hooks of length l as possible from Y ¹. It is called the l -core of Y .

Proposition 2.27. (i) If $k_i(Y) = 0$ for all $0 \leq i \leq l-1$, then $Y^{(l)} = \emptyset$.
(ii) There is a bijection between $\mathcal{C}^{(l)}$ and the set

$$\mathcal{K}^{(l)} \stackrel{\text{def.}}{=} \left\{ (k_0, \dots, k_{l-1}) \in \mathbb{Z}^l \mid \sum_{i=0}^{l-1} k_i = 0 \right\}.$$

Proof. The statement (i) follows from Remark 2.22(ii). To see (ii), it is enough to notice that $Y^{(l)}$ is uniquely determined by $(k_0(Y), \dots, k_{l-1}(Y))$ by Remark 2.20 and 2.22(ii). $\square$

Proposition 2.28 ([21] Theorem 2.7.30.). For any $l \geq 2$, a Young diagram Y is uniquely determined by its l -core $Y^{(l)}$ and l -quotient $\underline{Y}_{(l)}^*$. Thus for given $\lambda \in \mathcal{C}^{(l)}$, there is a bijection between the set

$$\mathcal{P}(n : \lambda) \stackrel{\text{def.}}{=} \{Y \in \mathcal{P}(n) \mid Y^{(l)} = \lambda\}$$

and $\mathcal{P}_l(n)$. Moreover, the weight of Y is given by

$$|Y| = |Y^{(l)}| + l|\underline{Y}_{(l)}^*|.$$

2.3.3. Weights of cores. Recall that we place a Young diagram Y on the (x, y) -plane, as shown in Figure 2.

Definition 2.29. For $\nu \in \mathbb{Z}$, we define $N_\nu(Y)$ to be the number of boxes on $y = x - \nu$.

Proposition 2.30. For $Y \in \mathcal{C}^{(l)}$ which is determined by $(k_0(Y), \dots, k_{l-1}(Y))$, we have

$$(5) \quad |Y| = \sum_{i=0}^{l-1} \left\{ \frac{1}{2} l k_i(Y)^2 + i k_i(Y) \right\}.$$

Proof. For $Y \in \mathcal{C}^{(l)}$, $|Y| = \sum_{i=0}^{l-1} \sum_{k \in \mathbb{Z}} N_{kl+i}(Y)$. By definition, we also have

$$N_{\mu l}(Y) = \begin{cases} \#\{\mathbf{m}_Y(\nu) \mid \mathbf{m}_Y(\nu) = 1, \nu < \mu l\} = \sum_i \#\{\mathbf{m}_{Y_i^*}(n) \mid \mathbf{m}_{Y_i^*}(n) = 1, n < \mu\}, & \text{if } \mu \leq 0, \\ \#\{\mathbf{m}_Y(\nu) \mid \mathbf{m}_Y(\nu) = 0, \nu \geq \mu l\} = \sum_i \#\{\mathbf{m}_{Y_i^*}(n) \mid \mathbf{m}_{Y_i^*}(n) = 0, n \geq \mu\}, & \text{if } \mu \geq 0. \end{cases}$$

¹It can be shown that $Y^{(l)}$ is well-defined, i.e. it does not depend on the choice of hooks to be removed.

Hence we obtain,

$$\begin{aligned} \sum_{\mu \in \mathbb{Z}} N_{\mu l}(Y) &= N_0(Y) + \sum_{\mu=1}^{\infty} (N_{\mu l}(Y) + N_{-\mu l}(Y)) \\ &= \sum_{k_i > 0} k_i(Y) + \sum_{n=1}^{\infty} \left\{ \sum_{k_i(Y) > n} (k_i(Y) - n) + \sum_{k_i(Y) < -n} (-k_i(Y) - n) \right\}. \end{aligned}$$

The right hand side of the above equation is equal to

$$\begin{aligned} &\sum_{k_i(Y) > 0} \frac{1}{2} k_i(Y) (k_i(Y) - 1) + \sum_{k_i(Y) < 0} \frac{1}{2} (-k_i(Y)) (-k_i(Y) - 1) + \sum_{k_i(Y) > 0} k_i(Y) \\ &= \frac{1}{2} \sum_{i=0}^{l-1} k_i(Y)^2 + \sum_{i=0}^{l-1} k_i(Y) \\ &= \frac{1}{2} \sum_{i=0}^{l-1} k_i(Y)^2, \end{aligned}$$

where we have used Remark 2.3.1(i). If we replace $k_n(Y)$ by $k_n(Y) - 1$ in the above discussion, we obtain

$$\sum_{\mu \in \mathbb{Z}} N_{\mu l+n}(Y) = \frac{1}{2} \sum_{i=0}^{l-1} k_i(Y)^2 + \sum_{i=n}^{l-1} k_i(Y).$$

Summing them up, we obtain the claim. $\square$

2.4. Quotients and cores for $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$. Let Γ be a cyclic group of order l . Let $\mathbf{w} = [W]$ be an isomorphism class of r -dimensional Γ -module. Let $W = \bigoplus_{i=1}^r \mathbb{C} \otimes \rho_i$ be a decomposition of W into 1-dimensional Γ -modules. We regard $(\rho_1, \dots, \rho_r)$ as an element of $(\mathbb{Z}/l\mathbb{Z})^r$, where we fix an order of them: $\rho_i \leq \rho_{i+1}$. We give the $(\rho_1, \dots, \rho_r)$ -coloring to $\mathcal{P}_r(n)$. Let us call it $\mathbf{w}$ -coloring. For fixed $\mathbf{w}$, $\mathcal{P}_r(n)$ has the following decomposition into disjoint sets:

$$(6) \quad \mathcal{P}_r(n) = \bigsqcup_{\mathbf{v}} \mathcal{P}_{\mathbf{w}}(\mathbf{v}),$$

where $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ is the subset of $\mathcal{P}_r(n)$ whose number of i -colored boxes is equal to v_i for $0 \leq i \leq l-1$, if we give the $\mathbf{w}$ -coloring to Young diagrams. Here $\mathbf{v} = (v_0, \dots, v_{l-1}) \in (\mathbb{Z}_{\geq 0})^l$ runs over all elements with $|\mathbf{v}| \stackrel{\text{def.}}{=} \sum_{i=0}^{l-1} v_i = n$.

Notation 2.31. Hereafter in this paper, we use the following assumptions and notations on l -dimensional row vector.

- (i) We understand that indices of l -dimensional row vectors are in $\mathbb{Z}/l\mathbb{Z}$.
- (ii) Let $\mathbf{e}_j \in \mathbb{Z}^l$ ($j = 0, \dots, l-1$) be the row vector whose j -th component is 1 and the others are 0.
- (iii) Let $\boldsymbol{\delta}$ be the row vector defined by $\boldsymbol{\delta} \stackrel{\text{def.}}{=} \sum_{j=0}^{l-1} \mathbf{e}_j$.

Let us take $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$. Then all boxes on the line $y = x - kl - i$ are colored by the same color $i + j$. Since v_i is equal to the number of i -colored boxes in Y , we have

$$(7) \quad v_i = \sum_{\mu \in \mathbb{Z}} N_{\mu l + i - j}(Y).$$

Proposition 2.32. All Young diagrams in $\mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$ have the same l -core.

Proof. For $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$, let $\{\mathbf{m}_Y(\nu)\}_{\nu=-\infty}^{\infty}$ be the Maya diagram corresponding to Y . For all $m \leq 0$, we have

$$N_m(Y) = \#\{\mathbf{m}_Y(\nu) \mid \mathbf{m}_Y(\nu) = 1, \nu < m\},$$

and for all $m \geq 0$, we have

$$N_m(Y) = \#\{\mathbf{m}_Y(\nu) \mid \mathbf{m}_Y(\nu) = 0, \nu \geq m\}.$$

Then we have

$$\begin{aligned} v_{i+j} - v_{i+j+1} &= \sum_{\mu \in \mathbb{Z}} N_{\mu l + i}(Y) - \sum_{k \in \mathbb{Z}} N_{\mu l + i + 1}(Y) \\ &= \sum_{\mu \in \mathbb{Z}} (N_{\mu l + i}(Y) - N_{\mu l + i + 1}(Y)) \\ &= \sum_{\mu \geq 0} (N_{\mu l + i}(Y) - N_{\mu l + i + 1}(Y)) - \sum_{\mu < 0} (N_{\mu l + i + 1}(Y) - N_{\mu l + i}(Y)) \\ &= \sum_{\mu \leq 0} (1 - \mathbf{m}_Y(\mu l + i)) - \sum_{\mu < 0} \mathbf{m}_Y(\mu l + i) \\ &= \#\{\mathbf{m}_Y(\mu l + i) = 0 \mid \mu \in \mathbb{Z}_{\geq 0}\} - \#\{\mathbf{m}_Y(\mu l + i) = 1 \mid \mu \in \mathbb{Z}_{< 0}\} \\ &= \#\{\mathbf{m}_{Y_i^*}(\mu) = 0 \mid \mu \in \mathbb{Z}_{\geq 0}\} - \#\{\mathbf{m}_{Y_i^*}(\mu) = 1 \mid \mu \in \mathbb{Z}_{< 0}\} \\ &= k_i(Y). \end{aligned}$$

Now our claim follows from Proposition 2.27(ii). $\square$

By definition, a hook of length nl has n i -colored boxes for all $0 \leq i \leq l - 1$. We have the following

Corollary 2.33. For each $0 \leq j \leq l - 1$,

$$\mathcal{P}_{\mathbf{e}_j}(n\delta) = \mathcal{P}(nl : \emptyset).$$

In particular $\mathcal{P}_{\mathbf{e}_j}(n\delta)$ is independent of j .

Proof. In the proof of Proposition 2.32, we obtained the formula $k_i(Y) = v_{i+j} - v_{i+j+1}$. Thus, for $Y \in \mathcal{P}_{\mathbf{e}_j}(n\delta)$, $k_i(Y) = 0$ for all $0 \leq i \leq l - 1$. This means that $Y^{(l)} = \emptyset$, by Proposition 2.27(i). $\square$

3. QUIVER VARIETIES AND TORUS ACTIONS

In this section, we introduce quiver varieties and torus actions on them. We follow [42] and [56] closely.

3.1. Quiver varieties.

3.1.1. *Framed moduli space of torsion free sheaves.* Let V, W be complex vector spaces with $\dim V = n$, $\dim W = r$. We consider

$$\mathbf{M}_r(n) \stackrel{\text{def.}}{=} (Q \otimes \text{Hom}(V, V)) \oplus \text{Hom}(W, V) \oplus \text{Hom}(V, W),$$

where Q is a 2-dimensional complex vector space. We define an action of $\text{GL}(V)$ on $\mathbf{M}_r(n)$ by

$$(B, i, j) \mapsto ((\text{id}_Q \otimes g)B(\text{id}_Q \otimes g)^{-1}, ig^{-1}, gj), \quad g \in \text{GL}(V).$$

We define a map $\mu : \mathbf{M}_r(n) \rightarrow \text{End}(V)$ by

$$\mu(B, i, j) \stackrel{\text{def.}}{=} [B \wedge B] + ij.$$

Note that $\mu^{-1}(0)$ is invariant under the action of $\text{GL}(V)$.

Definition 3.1. We say $(B, i, j) \in \mathbf{M}_r(n)$ is *stable* if the following condition is satisfied: If a subspace S of V contains $\text{Im}(i)$ and $B(S) \subset Q \otimes V$, then $S = V$.

We define

$$\mathcal{M}_r(n) \stackrel{\text{def.}}{=} \{(B, i, j) \in \mu^{-1}(0) \mid (B, i, j) \text{ is stable}\} / \text{GL}(V),$$

and

$$\begin{aligned} \mathcal{M}_r(n)_0 &\stackrel{\text{def.}}{=} \mu^{-1}(0) // \text{GL}(V) \\ &= \text{the set of closed } \text{GL}(V)\text{-orbits in } \mu^{-1}(0). \end{aligned}$$

These are *projective quotient* and *affine quotient* respectively, in the sense of geometric invariant theory (GIT). By the general theory of GIT, There is a proper morphism $\pi : \mathcal{M}_r(n) \rightarrow \mathcal{M}_r(n)_0$. The following theorem is proved in [44, §2].

Theorem 3.2. (i) $\mathcal{M}_r(n)$ is a nonsingular complex algebraic variety of dimension $2nr$ and $\pi : \mathcal{M}_r(n) \rightarrow \mathcal{M}_r(n)_0$ is a resolution of singularities.

(ii) $\mathcal{M}_r(n)$ is isomorphic to the moduli space of the pair (E, ϕ) where E is a torsion free sheaf over $\mathbb{CP}^2$ of rank $E = r$, $c_2(E) = n$ which is locally free in a neighbourhood of $l_\infty = \{[0 : z_1 : z_2]\}$ and ϕ is an isomorphism $E|_{l_\infty} \rightarrow \mathcal{O}_{\mathbb{CP}^2}^{\oplus r}$ (framing at infinity).

Remark 3.3. When $\dim(W) = 1$, any torsion free sheaf E of rank 1 with the above condition is a subsheaf of $\mathcal{O}_{\mathbb{CP}^2}$ such that $\text{Supp}(\mathcal{O}_{\mathbb{CP}^2}/E)$ is a 0-dimensional subscheme of $\mathbb{C}^2$, so we recover the Hilbert schemes of points on $\mathbb{C}^2$: $(\mathbb{C}^2)^{[n]} \cong \mathcal{M}_1(n)$. Therefore, $\mathcal{M}_r(n)$ is a higher rank generalization of the Hilbert scheme of points. Note that $\mathcal{M}_r(n)_0$ is isomorphic to the n -th symmetric product $S^n(\mathbb{C}^2)$ of $\mathbb{C}^2$ and the above morphism π coincides with the Hilbert-Chow morphism $\pi : (\mathbb{C}^2)^{[n]} \rightarrow S^n(\mathbb{C}^2)$ (see Appendix A.1).

3.1.2. *Quiver varieties.* Let Γ be a finite cyclic subgroup of $\text{SL}(2, \mathbb{C})$ of order l . Let Q be the 2-dimensional Γ -module defined by the inclusion $\Gamma \subset \text{SL}(2, \mathbb{C})$. For Γ -modules V and W , let

$$\mathbf{M}_W(V) \stackrel{\text{def.}}{=} (Q \otimes \text{Hom}_\Gamma(V, V)) \oplus \text{Hom}_\Gamma(W, V) \oplus \text{Hom}_\Gamma(V, W),$$

where $(\)_\Gamma$ means Γ -invariant part. We define $\text{GL}_\Gamma(V)$ -action on $\mathbf{M}_W(V)$, and the map $\mu : \mathbf{M}_W(V) \rightarrow \text{End}_\Gamma(V)$ as above. We also define the stability condition by the same condition as Definition 3.1, where subspace $S \subset V$ is replaced by Γ -submodule. Then we define $\mathcal{M}_w(\mathbf{v})$ and $\mathcal{M}_w(\mathbf{v})_0$ exactly the same as above, where $\mathbf{v}$ and $\mathbf{w}$ are isomorphism classes of V and W as Γ -module respectively. Let $\{R_i\}_{i=0}^{l-1}$ be the set of

all irreducible representations of Γ , where R_0 is the trivial representation. We can denote $V = \bigoplus_{i=0}^{l-1} V_i \otimes R_i$, $W = \bigoplus_{i=0}^{l-1} W_i \otimes R_i$. Then $\mathbf{v}$ and $\mathbf{w}$ can be regarded as row vectors $(\dim V_0, \dots, \dim V_{l-1})$ and $(\dim W_0, \dots, \dim W_{l-1})$ respectively. We call $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ and $\mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$ *quiver varieties*. The restriction of $\pi : \mathcal{M}_r(n) \rightarrow \mathcal{M}_r(n)_0$ gives a proper morphism $\pi : \mathcal{M}_{\mathbf{w}}(\mathbf{v}) \rightarrow \mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$.

Theorem 3.4. (i) $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is a nonsingular complex algebraic variety and $\pi : \mathcal{M}_{\mathbf{w}}(\mathbf{v}) \rightarrow \mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$ is a resolution of singularities. If $\mathcal{M}_{\mathbf{w}}(\mathbf{v}) \neq \emptyset$, its dimension is given by

$$(8) \quad \dim_{\mathbb{C}} \mathcal{M}_{\mathbf{w}}(\mathbf{v}) = \mathbf{v} \tilde{\mathbf{C}}_{\Gamma}^t \mathbf{v} + 2\mathbf{v}^t \mathbf{w},$$

where $\tilde{\mathbf{C}}_{\Gamma}$ is the affine Cartan matrix of type $A_{l-1}^{(1)}$ and ${}^t(\cdot)$ means the transposition.

(ii) $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is isomorphic to the moduli space parameterising the pair (E, ϕ) , where E is a Γ -equivariant torsion free sheaf with $H^1(\mathbb{CP}^2, E \otimes \mathcal{O}_{\mathbb{CP}^2}(-l_{\infty})) \cong V$ (as a Γ -module) and ϕ is a Γ -invariant trivialization $E|_{l_{\infty}} \rightarrow \mathcal{O}_{l_{\infty}} \otimes W$.

The statements (i) is due to Nakajima [36]. The statement (ii) is stated in [56, §2.3.]. Note that there is a natural Γ -action on $\mathcal{M}_r(n)$ induced by that on $\mathbb{CP}^2$, if we fix a lift of the Γ -action to $\mathcal{O}_{l_{\infty}}^{\oplus r}$ by $\mathbf{w}$. Then the above theorem tells us that Γ -fixed point set $\mathcal{M}_r(n)^{\Gamma}$ of $\mathcal{M}_r(n)$ have the following decomposition:

$$(9) \quad \mathcal{M}_r(n)^{\Gamma} = \bigsqcup_{\substack{\mathbf{v} \\ |\mathbf{v}|=n}} \mathcal{M}_{\mathbf{w}}(\mathbf{v}).$$

3.1.3. For the latter purpose, let us introduce the so-called *ALE spaces*.

Theorem 3.5 ([24, 18]). (i) $\mathcal{M}_{\mathbf{e}_0}(\boldsymbol{\delta})_0$ is isomorphic to the simple singularity $\mathbb{C}^2/\Gamma$. (ii) $\pi : \mathcal{M}_{\mathbf{e}_0}(\boldsymbol{\delta}) \rightarrow \mathcal{M}_{\mathbf{e}_0}(\boldsymbol{\delta})_0$ is the minimal resolution. Here we regard $\boldsymbol{\delta}$ as the isomorphism class of the regular representation of Γ . $\mathcal{M}_{\mathbf{e}_0}(\boldsymbol{\delta})$ is called *ALE space*².

This result was first obtained by Kronheimer [24], and rediscovered by Ito-Nakamura [18] and Ginzburg-Kapranov (unpublished) independently.

Remark 3.6. Results in §3.1.2 and §3.1.3 holds for any finite subgroup of $\mathrm{SL}(2, \mathbb{C})$, with a minor modification.

3.2. Torus actions and their fixed point set.

3.2.1. *Torus actions on framed moduli spaces.* Let T_Q (resp. T_W) be a maximal torus in $\mathrm{GL}(Q)$ (resp. $\mathrm{GL}(W)$), and let $T = T_Q \times T_W$. Note that $T_Q \cong (\mathbb{C}^*)^2$ and $T_W \cong (\mathbb{C}^*)^r$. Let us consider the following T -action on $\mathbf{M}_r(n)$:

$$(B_1, B_2, i, j) \mapsto (t_1 B_1, t_2 B_2, i e^{-1}, t_1 t_2 e j),$$

where $t = (t_1, t_2) \in T_Q$, $e = \mathrm{diag}(e_1, \dots, e_r) \in T_W$, and we denote $B = (B_1, B_2) \in Q \otimes \mathrm{Hom}(V, V)$. This induces a T -action on $\mathcal{M}_r(n)$ and $\mathcal{M}_r(n)_0$. Note that the morphism $\pi : \mathcal{M}_r(n) \rightarrow \mathcal{M}_r(n)_0$ is T -equivariant with respect to this action. This torus action is studied in [44]. It turns out that T -fixed points in $\mathcal{M}_r(n)$ are parameterized by $\mathcal{P}_r(n)$ and T -fixed points in $\mathcal{M}_r(n)_0$ consists of single point.

²ALE represents *asymptotically locally Euclidean*, since $\mathcal{M}_{\mathbf{e}_0}(\boldsymbol{\delta})$ admits a Riemannian metric which approximates the flat metric on $\mathbb{C}^2/\Gamma$ at infinity.

3.2.2. Torus actions on quiver varieties. We can regard T_Q (resp. T_W) be a maximal torus in $\mathrm{GL}_\Gamma(Q)$ (resp. $\mathrm{GL}_\Gamma(W)$). Thus we see that $T = T_Q \times T_W$ also acts on $\mathbf{M}_W(V)$ and this induces an action of T on quiver varieties $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ and $\mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$. We can study T -fixed points on quiver varieties by using results in [44].

Proposition 3.7. (i) $(E, \phi) \in \mathcal{M}_r(n)^\Gamma$ is fixed by the T -action if and only if E has a decomposition $E = I_1 \oplus \cdots \oplus I_r$ satisfying the following conditions for $\alpha = 1, \dots, r$.

(a) I_α is an ideal sheaf of 0-dimensional subscheme Z_α whose support is a Γ -orbits in $\mathbb{C}^2$.

(b) Under ϕ , $I_\alpha|_{l_\infty}$ is mapped to the α -th factor of $\mathcal{O}_{l_\infty} \otimes W$.

(c) I_α is fixed by the T_Q -action.

(ii) The T -fixed point set in $\mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$ consists of a single point.

Proof. This follows from the same argument as [44, Proposition 2.9]. $\square$

3.2.3. Parameterization of fixed points. It is instructive to compare (6) with (9).

Proposition 3.8. There is a one-to-one correspondence between the T -fixed point set of $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ and $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$.

Proof. This follows from Proposition 3.7 and the same argument as in [44, Proposition 2.9.(2)]. $\square$

3.2.4. We compare Proposition 3.8 with the work of Nakajima [35]. The results are not used later. In [35], an action of a certain 1-parameter subgroup of T on $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is considered. And in the case of $v_0 = w_0 = 0$, fixed points are parametrized by Young tableaux.

Definition 3.9. Let $\mu = (\mu_1, \dots, \mu_l)$ be an l -tuple of nonnegative integers.

(i) A μ -tableau of shape Y is a Young diagram Y whose boxes are numbered with the figures from 1 to l such that the cardinality of the boxes with figure k is μ_k .

(ii) A μ -tableau of shape Y is said to be *row-increasing* if the entries in each row increase strictly from the left to the right.

Proposition 3.10 ([35] Lemma 5.8.). When $v_0 = w_0 = 0$, T -fixed points on $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ are parametrized by row increasing μ -tableaux of shape $\lambda = (1^{w_1}, 2^{w_2}, \dots, (l-1)^{w_{l-1}})^3$, where $\mu_k \stackrel{\text{def.}}{=} v_n + \sum_{i \geq k} u_i$ ($k = 1, \dots, l-1$) and $\mu_l \stackrel{\text{def.}}{=} v_{l-1}$.

Proposition 3.11. When $v_0 = w_0 = 0$, there is a one-to-one correspondence between $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ and the set of row increasing μ -tableau of shape $\lambda = (\lambda_1, \dots, \lambda_r) \stackrel{\text{def.}}{=} (1^{w_1}, 2^{w_2}, \dots, (l-1)^{w_{l-1}})$.

Proof. Note that, in this case, $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ consists of tuples of l -cores. Each row in a μ -tableau of shape λ corresponds to a l -core. A correspondence is given as follows. For the i -th row of a μ -tableau of shape λ , we denote $\mathfrak{C}_i$ by the set of contents on it. The data $(k_0, \dots, k_{l-1})$ of a l -core is given as follows. For $1 \leq n \leq \lambda_i$,

$$k_{n-\lambda_i-1} = \begin{cases} 0 & (n \in \mathfrak{C}_i) \\ -1 & (n \notin \mathfrak{C}_i), \end{cases}$$

and for $\lambda_i < n \leq l$,

$$k_{n-\lambda_i-1} = \begin{cases} 1 & (n \in \mathfrak{C}_i) \\ 0 & (n \notin \mathfrak{C}_i). \end{cases}$$

³This means that λ is a partition with $w_i = \#\{j \mid \lambda_j = i\}$.

Then it is easy to see that this is a bijection. $\square$

3.2.5. Characters at fixed points.

Notation 3.12. (i) We denote by $e_1, \dots, e_r$ 1-dimensional T -modules given by $T \rightarrow \mathbb{C}^*; t = (t_1, t_2, e_1, \dots, e_r) \mapsto e_i$. Similar notations are used for t_1 and t_2 .
(ii) For $a, b \in \mathbb{Z}$, we set

$$\delta_{a,b}^{(l)} \stackrel{\text{def.}}{=} \begin{cases} 1 & \text{if } a \equiv b \pmod{l}, \\ 0 & \text{if } a \not\equiv b \pmod{l}. \end{cases}$$

Let $W = \oplus_{i=1}^r \mathbb{C} \otimes \rho_i$ be a decomposition of a representative W of $\mathbf{w}$ into 1-dimensional Γ -modules as in §2.3. Then we have the following

Theorem 3.13. Let x be a T -fixed point of $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ corresponding to $(Y_1, \dots, Y_r) \in \mathcal{P}_{\mathbf{w}}(\mathbf{v})$. Then the T -module structure of $T_x \mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is given by

$$(10) \quad T_x \mathcal{M}_{\mathbf{w}}(\mathbf{v}) = \sum_{\alpha, \beta=1}^r N_{e_\alpha, e_\beta}^\Gamma(t_1, t_2),$$

where

$$(11) \quad \begin{aligned} N_{e_\alpha, e_\beta}^\Gamma(t_1, t_2) = & e_\beta e_\alpha^{-1} \times \\ & \left\{ \sum_{s \in Y_\alpha} \left(t_1^{-l_{Y_\beta}(s)} t_2^{a_{Y_\alpha}(s)+1} \right) \delta_{h_{Y_\beta, Y_\alpha}(s), \rho_\alpha - \rho_\beta}^{(l)} \right. \\ & \left. + \sum_{t \in Y_\beta} \left(t_1^{l_{Y_\alpha}(t)+1} t_2^{-a_{Y_\beta}(t)} \right) \delta_{h_{Y_\alpha, Y_\beta}(t), \rho_\beta - \rho_\alpha}^{(l)} \right\}. \end{aligned}$$

Proof. This follows by taking Γ -invariant part of the T -module given in [44, Theorem 2.11]. $\square$

4. ENUMERATIVE GEOMETRY ON QUIVER VARIETIES

This section is devoted to computations of some global invariants of quiver varieties. In §4.1, we compute the ‘equivariant volume’ of $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$. This is a simple application of Theorem 3.13 and localization theorem. The main part in this section is §4.2, where we consider generating functions of Poincaré polynomials and Euler characteristics of quiver varieties. Our computation is based on combinatorial arguments over *cores* and *quotients* of Young diagrams in §2.

4.1. Equivariant integrals on $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$.

4.1.1. Localization. First, we review some basic facts about equivariant integrals and the localization formula. Let M be an algebraic variety with an action of a torus T and $H_*^T(M)$ be the equivariant homology⁴(with $\mathbb{Q}$ -coefficient). The ring $H_*^T(M)$ is a module over $H_*^T(pt)$. We consider the T -equivariant integral $\int_M 1$. This integral takes values in the quotient field of $H_*^T(pt)$. We refer the reader [45, §4.1.] for a precise definition of the

⁴See [45, Appendix C] for a precise definition.

	I	II	III	IV	V
Y_1			$\emptyset$	$\emptyset$	
Y_2	$\emptyset$	$\emptyset$			

TABLE 1.

The elements of $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ for $\mathbf{w} = (1, 1)$ and $\mathbf{v} = (1, 1)$.

integral. Assume that M is smooth and T -fixed point set M^T is finite. Then the classical Atiyah-Bott localization theorem [1] says that

$$\int_M 1 = \sum_{x \in M^T} \frac{1}{e_x},$$

where e_x is the T -equivariant Euler class of $T_x M$. ($e_x \neq 0$ since x is isolated.)

4.1.2. We apply the localization formula to $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$. In this case, the integral is defined by equivariant pushforward of the fundamental class $[\mathcal{M}_{\mathbf{w}}(\mathbf{v})]$ to the unique T -fixed point o in $\mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$ by $\pi : \mathcal{M}_{\mathbf{w}}(\mathbf{v}) \rightarrow \mathcal{M}_{\mathbf{w}}(\mathbf{v})_0$ (see Proposition 3.7 (ii)). Furthermore equivariant Euler classes at fixed points are given by Theorem 3.13. Let $(\epsilon_1, \epsilon_2, \vec{a})$ be a coordinate on the Lie algebra of T given by $t_1 = e^{\epsilon_1}, t_2 = e^{\epsilon_2}, e_{\alpha} = e^{a_{\alpha}}$, and $\vec{a} = (a_1, \dots, a_r)$.

Proposition 4.1. The T equivariant integral on $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is given by

$$\int_{\mathcal{M}_{\mathbf{w}}(\mathbf{v})} 1 = \sum_{\{Y_{\alpha}\}_{\alpha=1}^r \in \mathcal{P}_{\mathbf{w}}(\mathbf{v})} \frac{1}{\prod_{\alpha, \beta=1}^r n_{e_{\alpha} e_{\beta}}^{\Gamma}(\epsilon_1, \epsilon_2, \vec{a})},$$

where

$$\begin{aligned} n_{e_{\alpha} e_{\beta}}^{\Gamma}(\epsilon_1, \epsilon_2, \vec{a}) &= \prod_{\substack{s \in Y_{\alpha} \\ h_{Y_{\beta}} Y_{\alpha}(s) \equiv \rho_{\alpha} - \rho_{\beta} \pmod{l}}} (-l_{Y_{\beta}}(s) \epsilon_1 + (a_{Y_{\alpha}}(s) + 1) \epsilon_2 + a_{\beta} - a_{\alpha}) \\ &\times \prod_{\substack{t \in Y_{\beta} \\ h_{Y_{\alpha}} Y_{\beta}(t) \equiv \rho_{\beta} - \rho_{\alpha} \pmod{l}}} ((l_{Y_{\alpha}}(t) + 1) \epsilon_1 - a_{Y_{\beta}}(t) \epsilon_2 + a_{\beta} - a_{\alpha}). \end{aligned}$$

Example 4.2. In rank 1 cases, we have

$$(12) \quad \int_{\mathcal{M}_{\mathbf{e}_j}(\mathbf{v})} 1 = \frac{1}{n! l^n (\epsilon_1 \epsilon_2)^n},$$

where $n = |\underline{Y}_{(l)}^*|$ for $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$. A proof of this identity is given in Proposition A.5.

Example 4.3. Let us take $r = 2, l = 2, \mathbf{v} = (1, 1)$, and $\mathbf{w} = (1, 1)$. There are five elements in $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$, as shown in Table 1. Characters at these fixed points are given as

follows:

$$\begin{aligned}
\text{I.} \quad & t_1 t_2^{-1} + t_2^2 + e_1 e_2^{-1} t_2^{-1} + e_2 e_1^{-1} t_1 t_2^2 \\
\text{II.} \quad & t_1^{-1} t_2 + t_1^2 + e_2 e_1^{-1} t_1^2 t_2 + e_1 e_2^{-1} t_1^{-1} \\
\text{III.} \quad & t_2^2 + t_1 t_2^{-1} + e_2 e_1^{-1} t_2^{-1} + e_1 e_2^{-1} t_1 t_2^2 \\
\text{IV.} \quad & t_1^{-1} t_2 + t_1^2 + e_2 e_1^{-1} t_1^{-1} + e_1 e_2^{-1} t_1^2 t_2 \\
\text{V.} \quad & e_1 e_2^{-1} t_2 + e_2 e_1^{-1} t_1 + e_2 e_1^{-1} t_2 + e_1 e_2^{-1} t_1,
\end{aligned}$$

and we have

$$\int_{\mathcal{M}_{\mathbf{w}}(\mathbf{v})} 1 = \frac{4\epsilon_1^2 + 10\epsilon_1\epsilon_2 + 4\epsilon_2^2 - (a_1 - a_2)^2}{\epsilon_1\epsilon_2 E(1, -1, 2, 1) E(1, -1, 1, 2) E(-1, 1, 2, 1) E(-1, 1, 1, 2)},$$

where $E(x, y, z, w) \stackrel{\text{def.}}{=} xa_1 + ya_2 + z\epsilon_1 + w\epsilon_2$.

Remark 4.4. It seems interesting to study the generating function of equivariant integrals (see §5.1.2 and §A.3).

4.2. Poincaré polynomials and Euler characteristics of $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$.

4.2.1. *Poincaré polynomials for rank 1 case.* First, we consider Poincaré polynomials for rank 1 case. Let us consider the following generating function of Poincaré polynomials:

$$\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) \stackrel{\text{def.}}{=} \sum_{\substack{m \geq 0 \\ \mathbf{v} \in (\mathbb{Z}_{\geq 0})^l}} b_m(\mathcal{M}_{\mathbf{e}_j}(\mathbf{v})) \mathbf{t}^m \mathbf{q}^{|\mathbf{v}|} \prod_{i=0}^{l-1} \mathbf{r}_i^{v_i},$$

where $\vec{\mathbf{r}} = (\mathbf{r}_0, \dots, \mathbf{r}_{l-1})$ and b_m is the m -th Betti number. By the same argument as [45, §3.3.], we have

$$(13) \quad \mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) = \sum_{\mathbf{v} \in (\mathbb{Z}_{\geq 0})^l} \mathbf{q}^{|\mathbf{v}|} \prod_{i=0}^{l-1} \mathbf{r}_i^{v_i} \sum_{Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})} \mathbf{t}^{\#\{s \in Y \mid h(s) \equiv 0 \pmod{l}, l(s) > 0\}}.$$

Example 4.5. When $l = 3$, $\mathbf{v} = (1, 2, 1)$, $\mathbf{w} = (0, 1, 0)$,

$$\mathcal{P}_{\mathbf{w}}(\mathbf{v}) = \left\{ \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & \star & & \\ \hline & & \star & \\ \hline & & & \diamond \\ \hline \end{array} \right\}.$$

In the right hand of (13), $\star$'s are counted but $\diamond$ is not.

From the combinatorial discussions in §2, we have the following factorization formula for $\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$.

Theorem 4.6.

$$\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) = \mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) \mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}}),$$

where

$$\mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) \stackrel{\text{def.}}{=} \sum_{\substack{\mathbf{v} \in \mathbb{Z}_{\geq 0}^l \\ v_0 = \dots = v_{l-1}}} \mathbf{q}^{|\mathbf{v}|} \prod_{i=0}^{l-1} \mathbf{r}_i^{v_i} \sum_{Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})} \mathbf{t}^{\#\{s \in Y \mid h(s) \equiv 0 \pmod{l}, l(s) > 0\}},$$

and

$$\mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}}) \stackrel{\text{def.}}{=} \sum_{Y \in \mathcal{C}^{(l)}} \mathbf{q}^{|Y|} \prod_{i=0}^{l-1} \mathbf{r}_i^{\sum_{\mu \in \mathbb{Z}} N_{\mu l + i - j}(Y)}.$$

Proof. Note that $\mathcal{Z}^{\text{quot}}$ (resp. $\mathcal{Z}_{\mathbf{e}_j}^{\text{core}}$) is the contribution from l -quotients (resp. l -cores) of $\mathcal{P}_{\mathbf{w}}(\mathbf{v})$ corresponding to T -fixed points on $\mathcal{M}_{\mathbf{e}_j}(\mathbf{v})$. By Corollary 2.33, we have

$$(14) \quad \mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) = \sum_{n \geq 0} \mathbf{q}^{nl} \prod_{i=0}^{l-1} \mathbf{r}_i^n \sum_{Y \in \mathcal{P}(nl; \emptyset)} \mathbf{t}^{\#\{s \in Y \mid h(s) \equiv 0 \pmod{l}, l(s) > 0\}}.$$

Thus $\mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}})$ is independent of $\mathbf{e}_j$. Then our claim follows from equation (7) and the definition of quotients and cores. $\square$

Lemma 4.7.

$$\mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}_{\text{reg}}) = \prod_{m=1}^{\infty} \frac{1}{(1 - \mathbf{q}_{\text{reg}}^m \mathbf{t}^{2m-2})(1 - \mathbf{q}_{\text{reg}}^m \mathbf{t}^{2m})^{l-1}},$$

where $\mathbf{q}_{\text{reg}} \stackrel{\text{def.}}{=} \mathbf{q}^l \prod_{i=0}^{l-1} \mathbf{r}_i$.

Proof. We have

$$\begin{aligned} \mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}_{\text{reg}}) &= \sum_{n \geq 0} \mathbf{q}^{nl} \prod_{i=0}^{l-1} \mathbf{r}_i^n \sum_{Y \in \mathcal{P}(nl; \emptyset)} \mathbf{t}^{\#\{s \in Y \mid h(s) \equiv 0 \pmod{l}, l(s) > 0\}} \\ &= \sum_{n \geq 0} \mathbf{q}_{\text{reg}}^n \left\{ \sum_{m \geq 0} b_m(\mathcal{M}_{\mathbf{e}_0}(n\delta)) \mathbf{t}^m \right\}. \end{aligned}$$

Thus $\mathcal{Z}^{\text{quot}}(\mathbf{t}, \mathbf{q}_{\text{reg}})$ is the generating function of the Poincaré polynomials of $\mathcal{M}_{\mathbf{e}_0}(n\delta)$. It is known [26, 57] that $\mathcal{M}_{\mathbf{e}_0}(n\delta)$ is diffeomorphic to the Hilbert scheme of n points on the ALE space. By applying the Göttsche's formula (19) to the ALE space, we obtain the result. $\square$

Notation 4.8. We use the following definition of the Riemann theta function:

$$\Theta(\mathbf{u} | \mathbf{F}) \stackrel{\text{def.}}{=} \sum_{\mathbf{n} \in \mathbb{Z}^{l-1}} \exp[2\pi\sqrt{-1}(\mathbf{n}^t \mathbf{u} + \frac{1}{2} \mathbf{n} \mathbf{F}^t \mathbf{n})],$$

where $\mathbf{u}$ is a complex $(l-1)$ -dimensional row vector, ${}^t(\)$ means the transposition, and $\mathbf{F}$ is an $(l-1) \times (l-1)$ matrix.

Lemma 4.9. We have

$$\mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}}) = \Theta(\mathbf{u}_j | \mathbf{F}_{\Gamma}),$$

where

$$\mathbf{u}_j \stackrel{\text{def.}}{=} \frac{1}{2\pi\sqrt{-1}} (\log \mathbf{q} \mathbf{r}_{1+j}, \dots, \log \mathbf{q} \mathbf{r}_{l-1+j})$$

and

$$\mathbf{F}_{\Gamma} \stackrel{\text{def.}}{=} \frac{\log \mathbf{q}_{\text{reg}}}{2\pi\sqrt{-1}} \mathbf{C}_{\Gamma},$$

where $\mathbf{C}_{\Gamma}$ is the Cartan matrix of type A_{l-1} .

Proof. By Proposition 2.30 and its proof, we have

$$\begin{aligned}
\mathcal{Z}_{\mathbf{e}_j}^{\text{core}}(\mathbf{q}, \vec{\mathbf{r}}) &= \sum_{Y \in \mathcal{C}^{(l)}} \mathbf{q}^{|Y|} \prod_{i=0}^{l-1} \mathbf{r}_i^{\sum_{\mu \in \mathbb{Z}} N_{\mu l + i - j}(Y)} \\
&= \sum_{(k_0, \dots, k_{l-1}) \in \mathcal{K}^{(l)}} \mathbf{q}^{\frac{1}{2}l \sum_{i=0}^{l-1} k_i^2 + \sum_{i=0}^{l-1} i k_i} \mathbf{r}_j^{\frac{1}{2} \sum_{i=0}^{l-1} k_i^2} \prod_{\mu=1}^{l-1} \mathbf{r}_{j+\mu}^{\frac{1}{2} \sum_{i=0}^{l-1} k_i^2 + \sum_{i=\mu}^{l-1} k_i} \\
&= \sum_{(k_0, \dots, k_{l-1}) \in \mathcal{K}^{(l)}} \mathbf{q}_{\text{reg}}^{\frac{1}{2} \sum_{i=0}^{l-1} k_i^2} \prod_{\mu=1}^{l-1} (\mathbf{q} \mathbf{r}_{j+\mu})^{\sum_{i=\mu}^{l-1} k_i} \\
&= \sum_{(n_1, \dots, n_{l-1}) \in \mathbb{Z}^{l-1}} \prod_{\mu=1}^{l-1} \mathbf{q}_{\text{reg}}^{n_\mu(n_\mu - n_{\mu+1})} (\mathbf{q} \mathbf{r}_{j+\mu})^{n_\mu},
\end{aligned}$$

where $n_\mu \stackrel{\text{def.}}{=} \sum_{i=\mu}^{l-1} k_i$. Comparing this equation with the theta function in Notation 4.8, we obtain the lemma. $\square$

Theorem 4.10. We have

$$\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \vec{\mathbf{r}}) = \frac{\Theta(\mathbf{u}_j | \mathbf{F}_\Gamma)}{\prod_{m=1}^{\infty} (1 - \mathbf{q}_{\text{reg}}^m \mathbf{t}^{2m-2})(1 - \mathbf{q}_{\text{reg}}^m \mathbf{t}^{2m})^{l-1}}.$$

Proof. This follows from Theorem 4.6, Lemmas 4.7 and 4.9. $\square$

Remark 4.11. As we mentioned in the introduction, Theorem 4.10 is obtained from the result of Nakajima [42, §5.2.] (see comments in §4.2.3 below). This was informed us by H. Nakajima after the original version of this paper was submitted to e-print archives. He also pointed out that our argument has a close parallel to a geometric Frenkel-Kac construction of the Fock spaces of affine Lie algebras due to Grojnowski [15] (see also [39, §9.5.]). The authors are grateful to him for these comments.

4.2.2. Euler characteristics for higher rank case. Let $\dim W = r$. We consider the following generating function of the Euler characteristics $e(\mathcal{M}_{\mathbf{w}}(\mathbf{v}))$ of $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$:

$$\mathcal{E}_{\mathbf{w}}(\mathbf{q}) \stackrel{\text{def.}}{=} \sum_{\mathbf{v}} e(\mathcal{M}_{\mathbf{w}}(\mathbf{v})) \mathbf{q}^{|\mathbf{v}|} \prod_{i=0}^{l-1} \mathbf{r}_i^{v_i}.$$

Corollary 4.12.

$$\mathcal{E}_{\mathbf{w}}(\mathbf{q}) = \mathbf{q}_{\text{reg}}^{\frac{rl}{24}} \frac{\prod_{j=0}^{l-1} \Theta(\mathbf{u}_j | \mathbf{F}_\Gamma)^{w_j}}{\eta(\mathbf{q}_{\text{reg}})^{rl}},$$

where $\eta(q) \stackrel{\text{def.}}{=} q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$ is the Dedekind eta function.

Proof. We have

$$\mathcal{E}_{\mathbf{w}}(\mathbf{q}) = \prod_{j=0}^{l-1} (\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t} = 1, \mathbf{q}, \vec{\mathbf{r}}))^{w_j}.$$

Then by Theorem 4.10, we have the claim. $\square$

4.2.3. Let us make some comments on representation theoretical aspects of the result in Theorem 4.10. By Nakajima's pioneering work, it is known that quiver varieties are deeply related to representations of (quantum) affine algebras $\mathfrak{g}$ (of type $A_{l-1}^{(1)}$ in our case). It is shown in [38, 40] that their (K -)homology groups have structures of representations of affine algebras. Roughly speaking, $\oplus_{\mathbf{v}} H_*(\mathcal{M}_{\mathbf{w}}(\mathbf{v}))$ is a direct sum of certain highest weight representations of $\mathfrak{g}$ determined by $\mathbf{w}$. When $\mathbf{w}$ is the class of trivial 1-dimensional Γ -module, this is the so-called *Fock space* representation of $\mathfrak{g}$. This is explained in [42, §5.2.]. From this view point, the generating function of Euler characteristics is the character of the representation and the generating function of Poincaré polynomials is the so-called *q,t-character* [41], which plays a fundamental role in representation theory.

5. RELATIONS TO INSTANTON COUNTING

In this section, we summarize recent developments of *instanton counting* in physics. In §5.1, we give a brief explanation of instanton counting, which is one of our motivations of the first part of this paper. In §5.2, we explain a part of results of Maeda *et al.* [32, 33, 34], where quotients and cores for Young diagrams appeared.

5.1. Instanton counting. Mathematically speaking, instanton counting means computations of global invariants on the instanton moduli space, such as the Euler characteristic and the volume of the moduli space. What kind of invariants one want to compute depends on what kind of physical theory one considers. As we mentioned in introduction, $\mathcal{M}_{\mathbf{w}}(\mathbf{v})$ is a resolution of the moduli space of instantons on $\mathbb{C}^2/\Gamma$, therefore our results in §4 can be regarded as an instanton counting on $\mathbb{C}^2/\Gamma$.

5.1.1. In four-dimensional (4D) topologically twisted $\mathcal{N} = 4$ supersymmetric (SUSY) Yang-Mills (YM) theory, the partition function is given by the generating function of *Euler characteristics* of instanton moduli spaces, as we considered in §4.2. Such a theory is studied extensively by Vafa and Witten [55]. See also [22, 52], which considered issues similar to ours.

5.1.2. Nekrasov [47] introduced the following generating function of *equivariant volumes* of the instanton moduli spaces, in the sense of §4.1, in studies of 4D $\mathcal{N} = 2$ SUSY YM theory:

$$(15) \quad Z^{\text{inst}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda) \stackrel{\text{def.}}{=} \sum_{n=0}^{\infty} \Lambda^{2nr} \int_{\mathcal{M}_r(n)} 1 .$$

Such a generating function is called (the instanton part of) Nekrasov's partition function (see also [5, 9]). Nekrasov conjectured that $F^{\text{inst}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda) \stackrel{\text{def.}}{=} \epsilon_1 \epsilon_2 \log Z^{\text{inst}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda)$ is regular at $\epsilon_1 = \epsilon_2 = 0$, and $F^{\text{inst}}(0, 0, \vec{a}; \Lambda)$ coincides with (the instanton part of) the celebrated Seiberg-Witten prepotential. This conjecture is proved independently by Nakajima and Yoshioka [44, 45], Nekrasov and Okounkov [48], and Braverman and Etingof [3, 4]. We do not give a detail about Seiberg-Witten theory here. We refer the mathematical oriented readers [45, §2] for the Seiberg-Witten prepotential.

5.1.3. There is a 5D version $Z_{5D}^{\text{inst}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda)$ of the Nekrasov's partition function (15), which was also introduced by Nekrasov [47]. Physically speaking, that is $\mathcal{N} = 1$ SUSY YM theory on $\mathbb{R}^4 \times S^1$ (in a certain supergravity background). Mathematically speaking, this means that we consider equivariant integrals in (15) in the sense of equivariant K -theory [46].

5.2. **Cores and perturbative gauge theory.** In this section, we consider 5D version of the Nekrasov's partition function and discuss a combinatorial aspect of it. We explain works of Maeda *et al.* [32, 33, 34], where quotients and cores for Young diagrams appeared in a study of 5D $\mathcal{N} = 1$ SUSY YM theory explained in §5.1.3.

5.2.1. So far, we consider only the instanton part of the partition function. There is the so-called perturbative part $Z_{5D}^{\text{pert}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda)$ of the partition function, which corresponds to the perturbative part of the Seiberg-Witten prepotential. Although we do not give a definition here, it is defined by an explicit formula (see, for example, [45, Appendix E.]). Then the full partition function is define by

$$(16) \quad Z_{5D}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda) = Z_{5D}^{\text{pert}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda) Z_{5D}^{\text{inst}}(\epsilon_1, \epsilon_2, \vec{a}; \Lambda).$$

5.2.2. *Results of Maeda et al.* Recall that T -fixed points on $\mathcal{M}_r(n)$ are parametrized by $\mathcal{P}_r(n)$ and an equivariant integral (in the sense of K -theory) over $\mathcal{M}_r(n)$ can be expressed as a sum over $\mathcal{P}_r(n)$ by localization theorem, as we discussed in §4.1. The authors of *loc.cite.* found the following identity:

$$(17) \quad Z_{5D}^{r=1}(\hbar, -\hbar; \Lambda) = \sum_{\vec{k} \in \mathcal{K}^{(N)}} Z_{5D}^{r=N}(\hbar, -\hbar, \vec{k}; \Lambda),$$

where the LHS of (17) is the partition function for $r = 1$ under the condition⁵

$$(18) \quad \epsilon_1 = -\epsilon_2 = \hbar$$

and summands in the RHS of (17) are the partition function for $r = N(\geq 2)$ under the condition (18) and specialization $\vec{a} = \vec{k}$, where $\vec{k} = (\tilde{k}_1, \dots, \tilde{k}_N)$ is given by

$$\tilde{k}_m = \hbar \left\{ k_m + \frac{1}{N} \left(m - \frac{N+1}{2} \right) \right\}$$

for $\vec{k} = (k_1, \dots, k_N) \in \mathcal{K}^{(N)}$. In *loc.cite.*, this identity was proved by the following way. First, $Z_{5D}(\hbar, -\hbar, \vec{k}; \Lambda)$ can be identified with the partition function of a statistical model of plane partitions (= 3D Young diagrams) whose r -core of the ‘main diagonal’ is $Y^{(r)}(\vec{k})$, which can be described by free fermions. Then, under this identification, the perturbative part $Z_{5D}^{\text{pert}}(\hbar, -\hbar, \vec{k}; \Lambda)$ is identified with the contribution of ground states, which correspond to N -cores $Y^{(N)}(\vec{k}) \in \mathcal{C}^{(N)}$ determined by $\vec{k} \in \mathcal{K}^{(N)}$. And the instanton part $Z_{5D}^{\text{inst}}(\hbar, -\hbar, \vec{k}; \Lambda)$ is identified with the contribution of excitations, which correspond to N -quotients. Notice that, by Proposition 2.28, we recover the set of all Young diagrams $\mathcal{P}$ if we vary all $\vec{k} = (k_1, \dots, k_N) \in \mathcal{K}^{(N)}$ and $\underline{Y} \in \mathcal{P}_N$. Taking these into account, we can see that the identity (17) holds, after some Schur function calculus.

⁵Under this condition, Nekrasov's partition function can be expressed in terms of special values of Schur functions.

It is worth mentioning that we have a similar factorization of the generating function $\mathcal{Z}_{\mathbf{e}_j}(\mathbf{t}, \mathbf{q}, \bar{\mathbf{r}})$ of Poincaré polynomials in rank 1 case into ‘core-part’ $\mathcal{Z}^{\text{core}}$ and ‘quotient-part’ $\mathcal{Z}^{\text{quot}}$ in Theorem 4.6. Thus combinatorial structures in our case and the case studied by Maeda *et al.* are essentially the same.

Remark 5.1. Under the condition (18), it is known that Nekrasov’s partition function has connections with topological strings [7, 20, 54, 58] and 2D YM theory [31]. See Remark A.6 for a comment on cases with $\epsilon_1 + \epsilon_2 \neq 0$.

APPENDIX A. HILBERT SCHEMES OF POINTS ON SURFACES

This appendix has two purposes. The one is to summarize some facts about the Hilbert schemes of points⁶ (§A.1). The other is to give another method to compute equivariant volumes of quiver varieties in rank 1 case (§A.2, A.3). We get a closed formula for the generating function of such quantities. By comparing with the result in §4.1, we find an interesting combinatorial identity (Proposition A.5).

A.1. Hilbert schemes of points.

A.1.1. Let X be a smooth quasi-projective surface. The Hilbert scheme $X^{[n]}$ of n points on X is, by definition, a smooth algebraic variety of dimension $2n$, which parametrizes the set of 0-dimensional subschemes in X with colength n . Topology of the Hilbert scheme $X^{[n]}$ is well-known. The generating function of the Poincaré polynomials is given by the following Göttsche’s formula [14]:

$$(19) \quad \sum_{n \geq 0, i \geq 0} b_i(X^{[n]}) \mathbf{t}^i \mathbf{q}^n = \prod_{m \geq 1, i \geq 0} (1 - (-1)^i \mathbf{t}^{2m-2+i} \mathbf{q}^m)^{(-1)^{i+1} b_i(X)}.$$

A.1.2. There exists a proper surjective morphism $\pi : X^{[n]} \rightarrow S^n X$, where $S^n X$ is the n -th symmetric product of X . This morphism, so called the *Hilbert-Chow morphism*, gives a resolution of singularities of $S^n X$. Note that if X admits a torus action, then there is an induced torus action on $X^{[n]}$ and the Hilbert-Chow morphism π is equivariant.

A.2. Torus actions on Hilbert schemes. We study torus actions on the Hilbert schemes of points on toric surfaces. All the results in this section are due to Ellingsrud-Strømme [8] (see also [28, 29, 51], for a recent progress).

A.2.1. *Toric surfaces.* Let N be a 2-dimensional lattice and Σ be a fan in $N_{\mathbb{R}}$. From these datum, we can construct an algebraic surface $X = X(N, \Sigma)$, equipped with an action of the 2-dimensional algebraic torus $T = (\mathbb{C}^*)^2$ with a dense open orbit. These surfaces are called toric surfaces, and it is known that they are normal and quasi-projective (see, for example, [13, 49] for detail).

A.2.2. *Torus actions on toric surfaces.* Let X be a smooth toric surface associated to (N, Σ) . Then the 2-dimensional torus $T = N \otimes_{\mathbb{Z}} \mathbb{C}^*$ acts on X . The T -fixed points correspond to vertices of 2-dimensional cones in Σ . In particular, the number of T -fixed points is equal to the Euler number $e(X)$ of X . Let $X^T = \{p_1, \dots, p_{e(X)}\}$. Then, by construction, there is an affine chart $U_i = \text{Spec } \mathbb{C}[x_i, y_i]$ around each p_i which is T -invariant. We call (x_i, y_i) a toric coordinate around p_i . We denote $w_{x_i}(t_1, t_2)$ (resp. $w_{y_i}(t_1, t_2)$) the weight of the T -action on x_i (resp. y_i).

⁶A basic reference on this subject is [39].

A.2.3. Torus actions on Hilbert schemes. As we noticed in §A.1.2, T acts also on $X^{[n]}$. First, we identify T -fixed points on $X^{[n]}$.

Lemma A.1. There is a one-to-one correspondence between T -fixed point set on $X^{[n]}$ and $\mathcal{P}_{e(X)}(n)$.

Proof. A 0-dimensional subscheme Z of length n on X is T -fixed if and only if Z is supported on T -fixed points $\{p_i\}$ on X . Each connected component Z_i of Z with $\text{Supp}(Z_i) = \{p_i\}$ defines a T -fixed point in $(U_i)^{[n_i]} \cong (\mathbb{C}^2)^{[n_i]}$ for some $n_i (\geq 0)$ with $\sum_{i=1}^{e(X)} n_i = n$. It is well-known that there is a one-to-one correspondence between T -fixed points in $(\mathbb{C}^2)^{[n]}$ and partitions of n (see [39], Chapter 5). This gives the assertion. $\square$

Lemma A.2. The weight decomposition of the cotangent space of $X^{[n]}$ at a T -fixed point corresponding to $\underline{Y} = (Y_1, \dots, Y_{e(X)}) \in \mathcal{P}_{e(X)}(n)$ is given by

$$\sum_{i=1}^{e(X)} \sum_{s \in Y_i} \left\{ w_{x_i}(t_1, t_2)^{l(s)+1} w_{y_i}(t_1, t_2)^{-a(s)} + w_{x_i}(t_1, t_2)^{-l(s)} w_{y_i}(t_1, t_2)^{a(s)+1} \right\}.$$

Proof. This is obvious from the classical result of Ellingsrud and Strømme [8] on the weight decomposition of the tangent space of $(\mathbb{C}^2)^{[n]}$ at T -fixed point (see [39] Proposition 5.8). $\square$

For a Young diagram Y , we set

$$n_Y(\eta_1, \eta_2) \stackrel{\text{def.}}{=} \prod_{s \in Y} \{(l(s) + 1) \eta_1 - a(s) \eta_2\} \{-l(s) \eta_1 + (a(s) + 1) \eta_2\}.$$

Here η_1 , and η_2 are parameters. Then we have the following

Proposition A.3. T -equivariant integral on $X^{[n]}$ is given by

$$(20) \quad \int_{X^{[n]}} 1 = \sum_{\underline{Y} \in \mathcal{P}_{e(X)}(n)} \prod_{i=1}^{e(X)} \frac{1}{n_{Y_i}(\log w_{x_i}(e^{\epsilon_1}, e^{\epsilon_2}), \log w_{y_i}(e^{\epsilon_1}, e^{\epsilon_2}))}.$$

Here we write $t_i = e^{\epsilon_i}$, where $\epsilon_i \in \text{Lie}(T)$.

A.2.4. We consider the following generating function (15):

$$Z_X(\epsilon_1, \epsilon_2; \Lambda) \stackrel{\text{def.}}{=} \sum_{n=0}^{\infty} \Lambda^{2n} \int_{X^{[n]}} 1.$$

Let us consider the case $X = \mathbb{C}^2$ equipped with a T -action defined by

$$(21) \quad (x, y) \mapsto (t_1 x, t_2 y).$$

It is shown in [44, §4] that

$$(22) \quad Z_{\mathbb{C}^2}(\epsilon_1, \epsilon_2; \Lambda) = \exp \left(\frac{\Lambda^2}{\epsilon_1 \epsilon_2} \right).$$

Then we have

$$(23) \quad Z_X(\epsilon_1, \epsilon_2; \Lambda) = \sum_{n=0}^{\infty} \sum_{\underline{Y} \in \mathcal{P}_{e(X)}(n)} \frac{\Lambda^{2|\underline{Y}|}}{\prod_{i=1}^{e(X)} n_{Y_i} (\log w_{x_i}(e^{\epsilon_1}, e^{\epsilon_2}), \log w_{y_i}(e^{\epsilon_1}, e^{\epsilon_2}))}$$

$$(24) \quad = \prod_{i=1}^{e(X)} Z_{\mathbb{C}^2}(\log w_{x_i}(e^{\epsilon_1}, e^{\epsilon_2}), \log w_{y_i}(e^{\epsilon_1}, e^{\epsilon_2}); \Lambda)$$

$$(25) \quad = \exp \left\{ \Lambda^2 \sum_{i=1}^{e(X)} \frac{1}{(\log w_{x_i}(e^{\epsilon_1}, e^{\epsilon_2})) (\log w_{y_i}(e^{\epsilon_1}, e^{\epsilon_2}))} \right\}.$$

A.3. An example. We study a particular case when X is the ALE space of type A_{l-1} , i.e. $X = \mathcal{M}_{e_0}(\delta)$ and compare the result in §A.2.4 and the result in §4.1.

A.3.1. Recall that the simple singularity of type A_{l-1} is the quotient space $\mathbb{C}^2/\Gamma$, where the action of Γ on $\mathbb{C}^2$ is given by $(x, y) \mapsto (\zeta x, \zeta^{-1}y)$. Here ζ is a primitive l -th root of unity. Note that the above action of Γ on $\mathbb{C}^2$ commutes with the T -action (21) on $\mathbb{C}^2$. It follows that $\mathbb{C}^2/\Gamma$ is a toric singularity. It is well-known [13, 49] that it has the toric minimal resolution. In [18], it is shown that X is isomorphic to the toric minimal resolution, so that X has affine charts $U_i = \text{Spec } \mathbb{C}[x_i, y_i]$ ($i = 1, \dots, l$) defined by

$$(26) \quad x_i = \frac{a^i}{b^{l-i}}, \quad y_i = \frac{b^{l+1-i}}{a^{i-1}},$$

where (a, b) is a coordinate of $\mathbb{C}^2$ on which Γ acts by $(a, b) \mapsto (\zeta a, \zeta^{-1}b)$.

Remark A.4. It is easy to see that there is a bijection between the set of these l affine charts and the set $\mathcal{P}_{e_0}(\delta)$. This correspondence is pointed out in [19, Corollary 3.10], where $\mathcal{P}_{e_0}(\delta)$ is identified with the set of so-called Γ -clusters.

From equation (26), we have

$$(27) \quad \log w_{x_i}(\epsilon_1, \epsilon_2) = i\epsilon_1 - (l-i)\epsilon_2,$$

$$(28) \quad \log w_{y_i}(\epsilon_1, \epsilon_2) = -(i-1)\epsilon_1 + (l+1-i)\epsilon_2.$$

By substituting these into (25), we get

$$(29) \quad Z_X(\epsilon_1, \epsilon_2; \Lambda) = \exp \left(\frac{\Lambda^2}{l\epsilon_1\epsilon_2} \right).$$

The coefficient of Λ^{2n} is given by

$$(30) \quad \frac{1}{n!l^n(\epsilon_1\epsilon_2)^n},$$

which is nothing but the T -equivariant volume of the orbifold $S^n(\mathbb{C}^2/\Gamma)$. This is the desired result, since natural morphism $X^{[n]} \rightarrow S^n(\mathbb{C}^2/\Gamma)$, induced by $X \rightarrow \mathbb{C}^2/\Gamma$, is T -equivariant.

A.3.2. We have another resolution of singularities of $S^n(\mathbb{C}^2/\Gamma)$, which is given by quiver varieties [17, §7.2.3] (see also [57]). Note that by Proposition 2.28 and Proposition 2.32, total weight $|\underline{Y}_{(l)}^*|$ of l -quotient of $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$ is independent of Y . Let $n = |\underline{Y}_{(l)}^*|$ for $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$. Then by using Nakajima's *reflection functor* [38], it can be shown that $\mathcal{M}_{\mathbf{e}_j}(\mathbf{v})$ is a T -equivariant resolution of singularities of $S^n(\mathbb{C}^2/\Gamma)$. Thus equivariant integral $\int_{\mathcal{M}_{\mathbf{e}_j}(\mathbf{v})} 1$ also gives the equivariant volume of $S^n(\mathbb{C}^2/\Gamma)$. Therefore we have the following identity.

Proposition A.5.

$$\sum_{Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})} \prod_{\substack{s \in Y \\ h(s) \equiv 0 \pmod{l}}} \frac{1}{(-l(s)\epsilon_1 + (a(s) + 1)\epsilon_2)((l(s) + 1)\epsilon_1 - a(s)\epsilon_2)} = \frac{1}{n!l^n(\epsilon_1\epsilon_2)^n},$$

where $n = |\underline{Y}_{(l)}^*|$ for $Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})$.

Since the above identity is purely combinatorial, it should be possible to give a combinatorial proof. In the case of $\epsilon_1 = -\epsilon_2 = \hbar$, we have the following combinatorial proof.

$$\begin{aligned} \sum_{Y \in \mathcal{P}_{\mathbf{e}_j}(\mathbf{v})} \prod_{s \in \underline{Y}_{(l)}^*} \frac{1}{-(lh(s)\hbar)^2} &= \frac{1}{l^{2n}(-\hbar^2)^n} \sum_{\substack{(n_0, \dots, n_{l-1}) \in \mathbb{Z}_{\geq 0}^l \\ n_0 + \dots + n_{l-1} = n}} \prod_{i=0}^{l-1} \sum_{Y \in P(n_i)} \prod_{s \in Y} \frac{1}{\hbar(s)^2} \\ &= \frac{1}{l^{2n}(-\hbar^2)^n} \sum_{\substack{(n_0, \dots, n_{l-1}) \in \mathbb{Z}_{\geq 0}^l \\ n_0 + \dots + n_{l-1} = n}} \prod_{i=0}^{l-1} \frac{1}{n_i!} \\ &= \frac{1}{n!l^n(-\hbar^2)^n}. \end{aligned}$$

However, the authors do not have a combinatorial proof when $\epsilon_1 + \epsilon_2 \neq 0$.

Remark A.6. It is remarkable that in [16], geometry of the Hilbert scheme is applied to problems of enumerative combinatorics which is related to Macdonald polynomials. Note also that in a physical literature [2], relations between Macdonald polynomials and instanton counting in 5D SUSY YM theory (see §5.1.2) is pointed out. It seems interesting to study further in this direction.

REFERENCES

- [1] M.F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology **23** (1984), no. 1, 1-28.
- [2] H. Awata and H. Kanno, *Instanton counting, Macdonald functions and the moduli space of D-branes*, JHEP **0505**, 039 (2005) [arXiv:hep-th/0502061].
- [3] A. Braverman, *Instanton counting via affine Lie algebras I: Equivariant J-functions of (affine) flag manifolds and Whittaker vectors*, CRM Proc. Lecture Notes **38** (2004), 113–132.
- [4] A. Braverman and P. Etingof, *Instanton counting via affine Lie algebras. II: From Whittaker vectors to the Seiberg-Witten prepotential*, preprint, arXiv:math.AG/0409441.
- [5] U. Bruzzo, F. Fucito, J. F. Morales and A. Tanzini, *Multi-instanton calculus and equivariant cohomology*, JHEP **0305**, 054 (2003) [arXiv:hep-th/0211108].
- [6] E. Date, M. Jimbo, A. Kuniba, T. Miwa, M. Okado, *Paths, Maya diagrams and representations of $\widehat{\mathfrak{sl}}(r, \mathbb{C})$* , Integrable systems in quantum field theory and statistical mechanics, 149–191, Adv. Stud. Pure Math., **19**, Academic Press, Boston, MA, 1989.

- [7] T. Eguchi and H. Kanno, *Topological strings and Nekrasov's formulas*, JHEP **0312**, 006 (2003) [arXiv:hep-th/0310235].
- [8] G. Ellingsrud and S.A. Strømme, *On the homology of the Hilbert scheme of points in the plane*. Invent. Math. **87** (1987), no. 2, 343–352.
- [9] R. Flume and R. Poghossian, *An algorithm for the microscopic evaluation of the coefficients of the Seiberg-Witten prepotential*, Int. J. Mod. Phys. A **18**, 2541 (2003) [arXiv:hep-th/0208176].
- [10] I.B. Frenkel and A. Savage, *Bases of representations of type A affine Lie algebras via quiver varieties and statistical mechanics*, Int. Math. Res. Not. 2003, no. **28**, 1521–1547.
- [11] F. Fucito, J. F. Morales and R. Poghossian, *Multi instanton calculus on ALE spaces*, Nucl. Phys. B **703**, 518 (2004) [arXiv:hep-th/0406243].
- [12] S. Fujii and S. Minabe, *Poincaré polynomials of G-Hilbert schemes via Combinatorics*, preprint, 2005.
- [13] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, **131**. The William H. Roever Lectures in Geometry. Princeton University Press, Princeton, NJ, 1993.
- [14] L. Göttsche, *The Betti numbers of the Hilbert scheme of points on a smooth projective surface*, Math. Ann. **286** (1990), no. 1-3, 193–207.
- [15] I. Grojnowski, *Instantons and affine algebras. I. The Hilbert scheme and vertex operators*, Math. Res. Lett. **3** (1996), no. 2, 275–291.
- [16] M. Haiman, *t, q-Catalan numbers and the Hilbert scheme*, Discrete Math. **193** (1998), no. 1-3, 201–224.
- [17] ———, *Combinatorics, symmetric functions, and Hilbert schemes*, Current developments in mathematics, 2002, 39–111, Int. Press, Somerville, MA, 2003.
- [18] Y. Ito and I. Nakamura, *McKay correspondence and Hilbert schemes*, Proc. Japan Acad. Ser. **A 72** (1996), 135–138 ; *Hilbert schemes and simple singularities*, New trends in algebraic geometry, Proc. European Algebraic Geometry Conf. , Warwick 1996, Cambridge University Press, 1999, 151–233.
- [19] Y. Ito, *Special McKay correspondence*, Geometry of toric varieties, 213–225, Semin. Congr., **6**, Soc. Math. France, Paris, 2002.
- [20] A. Iqbal and A. K. Kashani-Poor, *Instanton counting and Chern-Simons theory*, Adv. Theor. Math. Phys. **7**, 457 (2004) [arXiv:hep-th/0212279].
- [21] G. James and A. Kerber, *The Representation Theory of the Symmetric Group*, Addison-Wesley, Reading, MA, 1981.
- [22] M. Jinzenji and T. Sasaki, *An approach to $N = 4$ ADE gauge theory on $K3$* , JHEP **0209**, 002 (2002) [arXiv:hep-th/0203179].
- [23] V.G. Kac, *Infinite-dimensional Lie algebras*, Third edition. Cambridge University Press, Cambridge, 1990.
- [24] P.B. Kronheimer, *The construction of ALE spaces as hyper-Kähler quotients*, J. Differential Geom. **29** (1989), no. 3, 665–683.
- [25] B. Kronheimer and H. Nakajima, *Yang-Mills instanton on ALE gravitational instantons*, Math. Ann. **288**, 263–307(1990).
- [26] A. Kuznetsov, *Quiver varieties and Hilbert schemes*, preprint, arXiv:math.AG/0111092.
- [27] B. Leclerc and H. Miyachi, *Some closed formulas for canonical bases of Fock spaces*, Represent. Theory **6** (2002), 290–312 (electronic).
- [28] W-P. Li, Z. Qin, W. Wang, *Hilbert schemes, integrable hierarchies, and Gromov-Witten theory*, Internat. Math. Res. Notices **40** (2004), 2085–2104.
- [29] ———, *The cohomology rings of Hilbert schemes via Jack polynomials*, CRM Proceedings and Lecture Notes, vol. **38** (2004), 249–258.
- [30] G.I. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995.
- [31] S. Matsuura and K. Ohta, *Localization on D-brane and gauge theory/Matrix model*, arXiv:hep-th/0504176.
- [32] T. Maeda, T. Nakatsu, K. Takasaki and T. Tamakoshi, *Five-dimensional supersymmetric Yang-Mills theories and random plane partitions*, JHEP **0503**, 056 (2005) [arXiv:hep-th/0412327].
- [33] ———, *Free fermion and Seiberg-Witten differential in random plane partitions*, Nucl. Phys. B **715**, 275 (2005) [arXiv:hep-th/0412329].

- [34] T. Maeda, T. Nakatsu, Y. Noma and T. Tamakoshi, *Gravitational quantum foam and supersymmetric gauge theories*, arXiv:hep-th/0505083.
- [35] H. Nakajima, *Homology of moduli spaces of instantons on ALE spaces. I*, J. Differential Geom. **40** (1994), no. 1, 105–127.
- [36] ———, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416.
- [37] ———, *Resolutions of moduli spaces of ideal instantons on R^4* , Topology, geometry and field theory, 129–136, World Sci. Publishing, River Edge, NJ, 1994.
- [38] ———, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. **3**, 515–560.
- [39] ———, *Lectures on Hilbert schemes of points on surfaces*, University Lecture Series **18**, American Mathematical Society, Providence, RI, 1999.
- [40] ———, *Quiver varieties and finite-dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238.
- [41] ———, *t -analogue of the q -characters of finite dimensional representations of quantum affine algebras*, Physics and combinatorics, 2000 (Nagoya), 196–219, World Sci. Publishing, River Edge, NJ, 2001.
- [42] ———, *Geometric construction of representations of affine algebras*, Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002), 423–438, Higher Ed. Press, Beijing, 2002.
- [43] ———, *Reflection functors for quiver varieties and Weyl group actions*, Math. Ann. **327** (2003), no. 4, 671–721.
- [44] H. Nakajima and K. Yoshioka, *Instanton counting on blowup. I*, preprint, arXiv:math.AG/0306198, to appear in Invent. Math..
- [45] ———, *Lectures on instanton counting*, CRM Proceedings and Lecture Notes, vol. **38** (2004), 31–101.
- [46] ———, *Instanton counting on blowup. II: K-theoretic partition function*, preprint, arXiv:math.AG/0505553.
- [47] N. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, Adv. Theor. Math. Phys. **7** (2003) 831–864, [arXiv:hep-th/0206161].
- [48] N. Nekrasov and A. Okounkov, *Seiberg-Witten theory and random partitions*, arXiv:hep-th/0306238.
- [49] T. Oda, *Convex Bodies and Algebraic Geometry. An introduction to the theory of toric varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), **15**. Springer-Verlag, Berlin, 1988.
- [50] J.B. Olsson, *Combinatorics and representations of finite groups*, Lecture Notes in Mathematics at the University of Essen, **20**. Universitat Essen, Fachbereich Mathematik, Essen, 1993.
- [51] Z. Qin, W. Wang, *Hilbert schemes of points on the minimal resolution and soliton equations*, preprint, arXiv:math.AG/0404540.
- [52] T. Sasaki, *Affine Lie algebras and S-duality of $N = 4$ super Yang-Mills theory for ADE gauge groups on $K3$* , arXiv:hep-th/0404194.
- [53] R.P. Stanley, *Enumerative combinatorics Vol. 2*, Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press, Cambridge, 1999.
- [54] Y. Tachikawa, *Five-dimensional Chern-Simons terms and Nekrasov’s instanton counting*, JHEP **0402**, 050 (2004) [arXiv:hep-th/0401184].
- [55] C. Vafa and E. Witten, *A Strong coupling test of S duality*, Nucl. Phys. B **431**, 3 (1994) [arXiv:hep-th/9408074].
- [56] M. Varagnolo and E. Vasserot, *On the K-theory of the cyclic quiver variety*, Internat. Math. Res. Notices 1999, no. **18**, 1005–1028.
- [57] W. Wang, *Hilbert schemes, wreath products, and the McKay correspondence*, preprint, arXiv:math.AG/0012104.
- [58] J. Zhou, *Curve counting and instanton counting*, preprint, arXiv:math.AG/0311237.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, NAGOYA, 464-8602, JAPAN
E-mail address: fujii@yukawa.kyoto-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, NAGOYA, 464-8602, JAPAN
E-mail address: minabe@yukawa.kyoto-u.ac.jp